	Name:		Patier	nt Name:					
			Patient Name:						
Address:				Address:					
			City: State: Zip:						
City:			DOB:/ Phone: ()						
hone:	()		Allerg	ies:					
	· /		** All	prescriptions a	are intende	d for prescril	bed pati	ent **	
MALE	E PERFORMANCE & TRT								
	Medication (must write Testosterone)	Concentrat	tion	Supplied	Directi	Directions		Refil	
	Cypionate PLUS	Cypionate 200m		10ml	INJ	ml week	dv		
ğ	(sesame oil)	Enanthate 20		□ include INJ I	kit -			□ nor	
era	Bi-blend (sesame oil)	Cypionate 180mg/ml Propionate 20mg/ml		10ml □ include INJ l	I IN I WAAKIV		dy	□ □ nor	
ţ.	(sesame oii)	□ 50mg □100mg □ 200mg		□ 30gm □ 60g	Oam				
nen	transdermal gel			□ 90gm	Apply _	gm QD		□ no	
Testosterone Replacement Therapy	Gonadorelin Acetate (in solution)	200mcg/ml		□ 5ml □ 2 x 5i		0.5ml SQ □ 2 □ 3			
epls	Clomiphene Citrate / Anastrozole	□ 30mg / 0.125mg		□ include INJ I Months	i i	kit: 31 gage – 1ml syringe) PO QD		□ noi	
e R	(capsule)	□ 60mg / 0.5mg						□ PF	
eron.	DHEA	□ 10mg □ 15mg			apply	/ gm qD			
oste	□ cream □ capsule □ troche	□ 20mg □ mg		□ 30 □ 60 □ 9	□ 1 PO	□ 1 PO qD			
Fest	Pregnenolone	□ 50mg □ 100n		□ 30 □ 60 □ 9	apply	□ apply gm qD			
	□ cream □ capsule □ troche	□ 250mg □ mg		30 00 0	□ 1 PO	□ 1 PO qD		□ PR	
	Combination (1) cream	□ YES □ NO		□ 30 □ 60 □ 9	90 □ apply	□ apply gm qD		□ □ PR	
	Medication (must write controlled)	Concentration		Supplied	Directi	ons		Refi	
a	Sermorelin	1000mcg/ml		12ml					
& Nandrolone Deconate		. ccameg,		□ include INJ kit	i 🗆 INJ C	□ INJ 0.3ml SQ QD Mon – Fri		□ PR	
Nandrolone Deconate	Deconate	200mg/ml		10ml □ include INJ kit INJml weekly			□ □ noi		
Nar	□ RC64-Cachexia	□ 12mg □ 28		□ Troches					
∞ర	□ 57mg □ Dye-free Capsule (12mg, 25mg)								
	□ Troc	□ 57mg						□ no	
5			ssolve 🗆	¼ □ ½ □ 1 troche				□ no	
		he: completely dis	ssolve □ le by mou	¼ □½ □1 trocheuth QD 2mL	INJ 0.1 - (ie QD 0.2 SQ prior to se		□ no □ □ no	
141	Bremelanotide (sq injection)	he: completely dis sule: take 1 capsul 10,000mcg/n	ssolve le by mou	¼ □½ □ 1 troche uth QD 2mL □ include INJ kit	INJ 0.1 - 0 activity, ma	D.2 SQ prior to saximum 3 WK or	· 8 MO	□ no □ □ no □ □ PR	
4	□ Caps	he: completely dis sule: take 1 capsu	ssolve le by mou	¼ □½ □1 trocheuth QD 2mL	INJ 0.1 - 0 activity, ma	ie QD 0.2 SQ prior to se	· 8 MO nutes	□ no □ no □ no □ PR	
PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin	he: completely dis sule: take 1 capsul 10,000mcg/n 2000iu / 1mo	ssolve le by mou	14	INJ 0.1 - (activity, ma 1 PO und- prior	D.2 SQ prior to so eximum 3 WK or er tongue 30 mir to sexual activity	nutes y ue 30	no no no no no no no no	
141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche)	he: completely dissule: take 1 capsul 10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 2	ssolve le by mount le by mount le by mount le le by mount le le by mount le by	14	INJ 0.1 - (activity, ma 1 PO und- prior	D.2 SQ prior to so eximum 3 WK or er tongue 30 mir to sexual activity PO under tongurior to sexual ac	nutes y ue 30 ctivity	no	
PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin	he: completely dis sule: take 1 capsul 10,000mcg/n 2000iu / 1mo	ssolve le by mount g 125iu nn	14	INJ 0.1 - (activity, ma 1 PO und prior 1½ □ 1 minutes p	D.2 SQ prior to so eximum 3 WK or er tongue 30 mir to sexual activity	nutes y ue 30 ctivity	no no no no PF	
★ PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche) Medication Sildenafil with Apomorphine 2mg □ oxytocin 125 units	he: completely dissule: take 1 capsul 10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 2 Concentratio 25mg 50mg 100mg	ssolve le by mou	14	INJ 0.1 - (activity, ma 1 PO und prior 1½ 1 minutes p	D.2 SQ prior to seximum 3 WK or er tongue 30 min to sexual activity PO under tongue for to sexual activity Quantity 90 60 30 10	nutes y ue 30 ctivity R 1 1 2	no no no price pri	
★ PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche) Medication Sildenafil with Apomorphine 2mg □ oxytocin 125 units Vardenafil with Apomorphine 2mg	he: completely dissule: take 1 capsul 10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 2 Concentration 25mg 50mg 100mg 5mg 10mg	ssolve	1/4 1/2 1 troche uth QD 2mL include INJ kit 10 20 30 10 20 30 Supplied Troches Dye-free Capsule Troches	INJ 0.1 - (activity, ma 1 PO und prior 1 ½ 1 minutes p	D.2 SQ prior to seximum 3 WK or er tongue 30 min to sexual activity. PO under tongue rior to sexual activity. Quantity. 90 60 30 10 90 60	nutes y ue 30 ctivity R 1 1 2 PRN 1 1 2	no no no no no pR	
★ PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche) Medication Sildenafil with Apomorphine 2mg oxytocin 125 units Vardenafil with Apomorphine 2mg oxytocin 125 units	10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 2 Concentratio 25mg 50m 100mg 5mg 10m 20mg	ssolve	14	INJ 0.1 - (activity, ma 1 PO und prior 1 ½ 1 minutes p	D.2 SQ prior to seximum 3 WK or er tongue 30 min to sexual activity PO under tongue for to sexual activity Quantity 90 60 30 10	nutes y ue 30 ctivity R 1 1 2 PRN 1 1 2	no no no no no no no no	
★ PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche) Medication Sildenafil with Apomorphine 2mg	he: completely discule: take 1 capsul 10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 2 Concentratio 25mg 50m 100mg 50mg 100 200mg 200mg	ssolve	1/4 1/2 1 troche Ith QD 2mL Include INJ kit 10 20 30 30 Troches 30 Troches 20 Troches	INJ 0.1 - (activity, ma 1 PO und- prior 1½	D.2 SQ prior to seximum 3 WK or er tongue 30 min to sexual activity PO under tongue rior to sexual ac Quantity 90 60 30 10 90 60 30 10	R MO nutes y ue 30 ctivity R	no no no no no no no no	
★ PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche) Medication Sildenafil with Apomorphine 2mg	10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 25mg 50mg 100mg 50mg 100mg 50mg 100mg 200mg 200mg 200mg 200mg 14 14 14 14 14 14 14 1	ssolve	1/4 1/2 1 troche Ith QD 2mL	INJ 0.1 - (activity, ma 1 PO und prior 1½	D.2 SQ prior to seximum 3 WK or er tongue 30 min to sexual activity PO under tongue rior to sexual ac Quantity 90 60 30 10 90 60 30 10	R MO nutes y ue 30 ctivity R	no no no no no no no no	
★ PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche) Medication Sildenafil with Apomorphine 2mg	10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 25mg 50mg 100mg 50mg 100mg 50mg 100mg 200mg 200mg 200mg 200mg 14 14 14 14 14 14 14 1	ssolve	1/4 1/2 1 troche Ith QD 2mL	INJ 0.1 - (activity, ma 1 PO und prior 1½	D.2 SQ prior to seximum 3 WK or er tongue 30 min to sexual activity PO under tongue rior to sexual ac Quantity 90 60 30 10 90 60 30 10	R MO nutes y ue 30 ctivity R	noi	
PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche) Medication Sildenafil with Apomorphine 2mg	10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 25mg 50mg 100mg 50mg 100mg 200mg 50mg 100mg 200mg 5mg 100mg 200mg 5mg 100mg 200mg 5mg 10mg 200mg 200mg 200mg 20mg 20	ssolve	1/4 1/2 1 troche Ith QD 2mL Include INJ kit 10 20 30 30 10 20 30 Supplied Troches Dye-free Capsule	INJ 0.1 - (activity, ma 1 PO und prior 1½ 11 minutes p	D.2 SQ prior to seximum 3 WK or er tongue 30 min to sexual activity PO under tongue for to sexual activity 90 60 30 10 90 60 30 10 90 60 30 10 10 10 10 10 10 10 10 10	* 8 MO nutes y ue 30 ctivity R	noi	
★ PT141	Bremelanotide (sq injection) Bremelanotide with Methylcobalamin (sublingual troche) Tadalafil / Bremelanotide / Oxytocin (sublingual troche) Medication Sildenafil with Apomorphine 2mg	10,000mcg/n 2000iu / 1mg 25mg / 2000iu / 25mg 50mg 100mg 50mg 100mg 50mg 100mg 200mg 5mg 10mg 200mg 5mg 10mg 20mg 5mg 10mg 20mg 5mg 10mg 20mg 5mg 10mg 20mg 10mg 20mg 10mg 20mg 10mg 20mg 10mg 20mg 10mg 20mg 10mg 10mg 20mg 10mg 10mg	ssolve	1/4 1/2 1 troche Ith QD	INJ 0.1 - (activity, ma 1 PO und prior 1½	D.2 SQ prior to seximum 3 WK or er tongue 30 min to sexual activity PO under tongue for to sexual activity 90 60	* 8 MO nutes y ue 30 ctivity R	noi	

Office: (855) 277-2488 Fax: (888) 689-9892